



Contribution ID: 134

Type: Poster

## Unconventional superconductivity in topological ruthenium silicides with Kramers and hourglass fermions

Monday, 29 August 2022 18:40 (20 minutes)

The convergence of two major research strands in modern condensed-matter physics: topological materials and unconventional superconductivity, constitutes a new field of study. Topological materials with Kramers or hourglass fermions represent a special subclass, recently realized in materials lacking inversion symmetry or with a nonsymmorphic space group. At the same time, there is a surge of interest in identifying time-reversal symmetry (TRS) breaking (a key feature of unconventional superconductivity) in this class of materials, as a new routine way to realize topological superconductivity.

By using the muon-spin rotation and relaxation technique, backed by detailed theoretical analyses, we show that TRuSi ( $T = \text{Ti, Nb, Hf, and Ta}$ ) noncentrosymmetric materials represent a family of compounds encompassing all the above unique properties [1]. Their bulk normal states behave as three-dimensional Kramers nodal-line semimetals, characterized by a fairly large antisymmetric spin-orbit coupling and by glide-reflection-protected hourglass-like fermions. We also identify surface states near the Fermi level of TRuSi materials. More interestingly, NbRuSi and TaRuSi undergo a superconducting transition, which spontaneously breaks TRS below  $T_c$ , while surprisingly showing a fully-gapped superconducting ground state. This superconducting ground state is consistent with a unitary ( $s + ip$ ) pairing, i.e., with a mixture of spin-singlet and spin-triplet pairings. As such, the TRuSi family provides an ideal platform for investigating the rich interplay between the exotic properties of Kramers nodal-line/hourglass fermions and unconventional superconductivity.

[1] T. Shang, J. Z. Zhao, et al., and T. Shiroka, submitted to *Sci. Adv.* (2022).

**Primary authors:** SHIROKA, T. (Laboratorium für Festkörperphysik, ETHZ); ZHAO, J. Z. (Co-Innovation Center for New Energetic Materials, Southwest University of Science and Technology,); HU, L.-H. (Department of Physics and Astronomy, University of Tennessee); MA, J. Z. (Department of Physics, City University of Hong Kong); GAWRYLUK, D. J. (Laboratory for Multiscale Materials Experiments, Paul Scherrer Institut); ZHU, X. Y. (Key Laboratory of Polar Materials and Devices (MOE), School of Physics and Electronic Science, East China Normal University); ZHANG, H. (Key Laboratory of Polar Materials and Devices (MOE), School of Physics and Electronic Science, East China Normal University); ZHEN, Z. X. (Key Laboratory of Polar Materials and Devices (MOE), School of Physics and Electronic Science, East China Normal University); YU, B. C. (Key Laboratory of Polar Materials and Devices (MOE), School of Physics and Electronic Science, East China Normal University); XU, Y. (Key Laboratory of Polar Materials and Devices (MOE), School of Physics and Electronic Science, East China Normal University); ZHAN, Q. F. (Key Laboratory of Polar Materials and Devices (MOE), School of Physics and Electronic Science, East China Normal University); POMJAKUSHINA, E. (Laboratory for Multiscale Materials Experiments, Paul Scherrer Institut); SHI, M. (Swiss Light Source, Paul Scherrer Institut); SHANG, T. (Key Laboratory of Polar Materials and Devices (MOE), School of Physics and Electronic Science, East China Normal University)

**Presenter:** SHIROKA, T. (Laboratorium für Festkörperphysik, ETHZ)

**Session Classification:** Posters

**Track Classification:** Strongly correlated electron systems